



VAGINAL SUPPOSITORY

Treatment of cervical erosion, vaginitis, vaginal abnormal discharge, and other related symptoms.


Vaginal suppository in KMUH

Polinin 90mg Vaginal Tablet
保理寧陰道栓劑



高雄醫學大學附設中和紀念醫院版權所有 109.03.11

Lomexin vaginal soft cap. 200mg
洛徽欣陰道軟膠囊 200毫克



Esvatin VT 0.5mg/Tab
薇詩婷陰道錠



高雄醫學大學附設中和紀念醫院版權所有 11

Before using vaginal suppository



Wash your vaginal area and hands

Wash with mild soap and warm water, and dry well with a clean towel.

How to use vaginal suppository

1 Remove any wrapping from the suppository.

If there is no wrap around, skip this step.

2 Lie on your back with your knees bent and legs slightly apart.



3 Gently insert the suppository into your vagina **with its pointed end first** and as far as it will comfortably go. (approximately two finger joints deep.)

4 It is best to take the medication **before bedtime** to prolong its duration and enhance its effectiveness





VAGINAL SUPPOSITORY

Be aware of...

- If you are allergic to medications, you should inform your doctor or pharmacist in advance.
- During the treatment, even if symptoms improve, do not discontinue the medication on your own; **complete the entire course for a complete cure.**
- During the infection treatment with medication, avoid sexual activities, or request males to use condoms to prevent reinfection.
- During infection treatment, avoid using irritating cleaning products.
- Due to different properties of drug products and patients' conditions, please ask your doctor or pharmacist whether or not to continue using the drug during your menstrual period.
- To prevent the suppository from dissolving and staining clothing, it is recommended to use a sanitary pad.

Pregnant or breastfeed

If planning to become or already pregnant, you should inform and discuss with your doctor, as well as before planning to breastfeed.

Storage

- Store in a **cool, dry place at room temperature.** Keep it out of the reach of children to avoid accidental ingestion.
- If the suppository becomes soft, you can place it in cold water or in the refrigerator. Use it after it getting hardens.

Possible side effects that may occur

- In the early stages of treatment, there may be localized irritation, but it typically subsides quickly.