



Loceryl[®] nail lacquer

(5% w/v amorolfine HCl)

For Onychomycosis




How to use Loceryl[®] nail lacquer




1

Before using Loceryl[®], file the infected area of the nails smoothly as much as possible.




2

Use a cleaning pad to clean the infected nail.




3

Dip the reusable spatula into the Loceryl[®] nail lacquer. Do not wipe it on the bottle edge.




4

Paint the Loceryl[®] nail lacquer to cover the entire infected nail.



5

After finishing treatment, use the cleaning pad to clean the spatula and the neck of the bottle.



6

Close the bottle tightly right to prevent it from drying out.



7

Allow the nail to dry for 3–5 minutes.

Where to apply the lacquer

1. Apply the Loceryl[®] nail lacquer to the affected fingernails or toenails once or twice a week.
2. Do not file the skin around the nail.



Loceryl[®] nail lacquer

Be aware of...

1. Treatment usually takes about **6 months for fingernails** and **9–12 months for toenails**.
2. Continue using Loceryl[®] until a new, healthy nail has fully grown.
3. **Do not stop treatment early**, as this may cause the infection to return. Nail fungus treatment **requires patience and regular use**. Stopping early may cause relapse.
4. Wear impermeable gloves if you work with organic solvents (e.g. thinners or mineral spirits).
5. If there is no improvement **after 3 months**, see your doctor for further evaluation.
6. **Maintaining good hygiene** (keep nails dry, do not share nail tools, shoes, or socks) helps improve treatment success.

Forget to use

- Apply the medicine as soon as you remember and continue as scheduled
- Do not use a double dose

Pregnancy

- This medicine is not recommended during pregnancy or breastfeeding

Storage

- Store below 30°C in a cool place
- Avoid sunlight

Possible side effects

- Mild burning or redness may occur
- Nails may become discolored or brittle