




Regaine[®] 5% Women's Foam

(Minoxidil 60g/Bot)

For patients with androgenetic alopecia



How to use Regaine[®]




- 1 Hold the ring and rotate it to align the white arrow on the ring with the gray arrow on the cap. Pull the cap back and up to open.
- 2 Hold the can **completely upside down** and dispense the foam onto dry fingers. The total amount used should be **half a capful (1g)**. (Tip: If your fingers are warm, the foam may melt. Cool your hands with cold water and dry them thoroughly before use.)
- 3 Part your hair in the thinning area to expose the scalp as much as possible. Use your other hand to apply the foam directly to the scalp in that area.
- 4 Push the cap down onto the can and ensure the arrows are **NOT aligned**. This prevents children from opening it easily.

Where and how to apply

1. Apply directly to the hair loss area on the scalp.
2. Use 1 g (half a capful) once a day.

WRONG DISPENSING (X marks error)





Regaine® 5% Women's Foam

Be aware of...

1. This medication is for topical use on the **scalp only**.
2. It usually takes **3 to 6 months** to see hair growth effects.
3. Temporary hair loss may occur in the **first 2 weeks** as follicles enter a new growth cycle. If shedding continues or worsens, **stop use and consult a doctor**.
4. Hair and scalp must be **completely dry** when applying the foam.
5. Avoid getting the scalp wet **for 4 hours** after use.
6. Remember to **wash your hands thoroughly after use**.
7. If the foam touches your eyes, wounds, or mucous membranes, **rinse immediately** with plenty of cold water.

Forget to use

If you forget, apply it as soon as you remember. If it is almost time for your next dose, skip the missed one. Do not use a double dose.

Pregnancy

If you are pregnant or breast-feeding, do not use this medication.

Storage

- Store at or below 30°C
- Keep until the expiration date on the carton.
- Keep away from fire, flame, or smoking during and after use.

Possible side effects

- Allergic rhinitis, shortness of breath, headache, blood pressure/heart rate changes, edema.
- If dermatitis or other side effects occur, stop use and tell your doctor.