



Regaine® 5% Extra Strength External Solution (Minoxidil 60mL/Bot)

For patients with androgenetic alopecia




How to use Regaine®

For Large Areas — Regular Spray Applicator



- 1 Remove the large outer black cap and the internal black bottle cap.
- 2 Insert the regular spray applicator into the bottle and screw it on tightly.
- 3 Aim at the hair loss area, press once, and use your fingertips to spread evenly. Repeat **6 times** in total to deliver a full 1 mL dose.

For Small Areas — Extended Spray Tip



- 1 The regular applicator base must be installed first.
- 2 Remove the regular spray head, attach the extended tip firmly, and open the small cap.
- 3 Aim at the hair loss area, press once, and use your fingertips to spread evenly. Repeat **6 times** in total to deliver a full 1 mL dose.

Where and how to apply

1. Apply directly to the hair loss area on the scalp.
2. Use 1 mL (6 sprays) twice a day (morning and evening).



Regaine® 5% Extra Strength External Solution

Be aware of...

1. This medication is for topical use on the **scalp only**.
2. It usually takes **2 to 4 months** to see hair growth effects.
3. Temporary hair loss may occur in the **first 2 weeks** as follicles enter a new growth cycle. If shedding continues or worsens, **stop use and consult a doctor**.
4. Hair and scalp must be **completely dry** when applying the solution.
5. Avoid getting the scalp wet **for 4 hours** after use.
6. Remember to **wash your hands thoroughly after use**.
7. If the solution touches your eyes, wounds, or mucous membranes, **rinse immediately** with plenty of cold water.

Forget to use

If you forget, apply it as soon as you remember. If it is almost time for your next dose, skip the missed one. Do not use a double dose.

Pregnancy

If you are pregnant or breast-feeding, do not use this medication.

Storage

- Store at room temperature. (15–25°C)
- Avoid sunlight.
- After opening, it can be used until the expired date.

Possible side effects

- Allergic rhinitis, shortness of breath, headache, blood pressure/heart rate changes, edema.
- If dermatitis or other side effects occur, stop use and tell your doctor.