




# Azetin<sup>®</sup> Nasal Spray

(Azelastine hydrochloride 140 mcg/Dose, 120 Dose/Bot)

For allergic rhinitis and nasal symptom



## Prepare for first time use



1 Remove the **safety ring** and **cap**

2 Spray several times to ensure the plenty amount of spray


## How to handle the device

Hold nasal spray upright, thumb beneath the bottle, fingers on either side of the nozzle.

## Using the device regularly



Created by Melissa Moller from Noxin Project



Created by Inez Tan Kuber from Noxin Project

1 Blow your nose gently.

2 Remove the cap.

3 Tilt your head forward slightly. Place the nozzle in one of your nostrils. Point the end of the nozzle toward the outside of your nose

4 Breathe in gently through your nose and press the applicator down firmly between your fingers and thumb.

5 Take the nozzle out and breathe out through your mouth.

6 Wipe the nozzle. Then replace the cap





# Azetin<sup>®</sup> Nasal Spray

## Be aware of...

1. Do not spray in the direction of the center ridge of nose
2. Do not rub the nozzle with any needle or sharp object
3. When spraying, keep the bottle upright
4. Do not give the medicine to your family or friends
5. Be careful when driving or operating dangerous machinery after using the Azetin.
6. Avoid to use alcohol simultaneously, due to exacerbate central depression effect.
7. Maybe produce a bitter taste

### Possible side effect

- Nasal mucosal irritation (e.g. tingling, itching, sneezing, even bleeding from the nose)
- Others such as headache, sleepiness, dizziness, dry mouth, dry nose, weight gain, myalgia, skin rash, anxiety, depression

### Forget to use

Whenever recalling a missing dose, take it immediately. Do not take the missing dose if next instructed dosing time is approaching

### How to store?

- Store below 25 degrees.
- Avoid sunlight
- It can be stored for two months after opening

