


VENTOLIN® INHALER

(Salbutamol 100 mcg /dose)

Treatment or prevention of bronchospasm



A modern delivery system



Before using a new inhaler for the first time



Prime the inhaler


- ① Take the cap off the mouthpiece and shake the inhaler well.
- ② Shake and spray the inhaler 3 more times into the air away from your face to finish priming it. Avoid spraying in eyes.

Prime your inhaler again if you have not used it in more than 14 days or if you drop it.

How to use VENTOLIN inhaler



Shake the inhaler well before each spray. Take the cap off the mouthpiece of the actuator.



Insert inhaler into spacer (if necessary).



Breathe out gently. Put the mouthpiece in your mouth and close your lips around it.




4 Hold the inhaler with the mouthpiece down.




VENTOLIN® INHALER

(Salbutamol 100 mcg /dose)





Hold your breath for 5~10 seconds.



Remove the inhaler and **put the cap back on the mouthpiece** after every time you use it. **Make sure it snaps firmly into place.**Cap

If your need to use <u>more sprays</u>, <u>wait 30 secs to 1 min</u> and <u>shake the inhaler again</u>.

Repeat Steps 1 through Step 6.

Be aware of...

Get medical help right away if VENTOLIN no longer helps your symptoms (ex: wheezing/trouble breathing), or you need to use your inhaler more often.

Attention !!!

This medication is for urgent condition. Following the direction from your doctor. **Do not** increase your dose or take extra doses casually.

Cleaning!!!

- 1. The actuator and mouthpiece can be cleaned by wet or dry tissue.
- The spacer can be washed under the faucet and run warm water through it. Let the spacer air-dry overnight.

Storage

- Store at room temperature
- Avoid sunlight
- After opening, it can be used until the expired date

How do I know when to replace the inhaler

- VENTOLIN INHALER provides 200 doses.
- You may record how many doses you use every day to calculate dose remain in the inhaler.