






Foster Nexthaler[®]

(Beclometasone 100mcg/Formoterol 6mcg, 120 Dose/Bot)
For patients with **asthma** or **chronic obstructive pulmonary disease (COPD)**




Instructions for use



Check the counter


- Before the inhaler is used for the first time, the counter should show **the number 120**. This is the number of doses in the inhaler.
- Each time you open the cover, you prepare 1 dose of medicine.

Prepare your dose: Wait to open the cover until you are ready to take your dose.




1 Open the cover fully.

- Slide the cover down to expose the mouthpiece.
- Your inhaler is now ready to use.
- Before inhaling breathe out as far as is comfortable.




2 Inhale your medicine.

- Lift your inhaler up to your mouth and place your lips around the mouthpiece.
- Take a **forceful and deep breath** through your mouth.
- You should **hear a "click"** when you take your dose. Keep your inhalation for 1 to 2 seconds.
- Remove your inhaler from your mouth. Hold your breath for 3 to 4 seconds at least.



3 Close the inhaler.

- Close the cover fully. Check that the dose counter has gone down by one
- If necessary, you may clean your inhaler after use with a dry tissue. Do not clean it with water.





Foster Nexthaler[®]

(Beclometasone 100mcg/Formoterol 6mcg, 120 Dose/Bot)

For patients with **asthma** or **chronic obstructive pulmonary disease (COPD)**



Be aware of...

1. Please gargle with water or brush your teeth after inhaling.
2. The number shown in the window dose not decrease on closing the cover if the patient has not inhaled through the inhaler.
3. If the Dose Counter Window showed "0" there are no doses left – dispose of your inhaler and get a new one.
4. Do not give the medicine to your family or friends.



Possible undesirable effects

- The most common is mild tremor at the beginning of treatment.
- Other common included pneumonia (in COPD patients)
- Headache, tachycardia, sinus bradycardia, angina pectoris, myocardial ischaemia, QT prolongation is uncommon.
- nasopharyngitis, oral candidiasis, dysphonia, throat irritation, irritability, cortisol free urine decreased, blood cortisol decreased, blood glucose increased is uncommon.
- If the following symptom, please notify your doctor immediately: rashes, urticaria, pruritus and erythema and oedema of the eye, face, lips and throat (angioedema).

Forget to use

Whenever recalling a missing dose, take it immediately. Do not take the missing dose if next instructed dosing time is approaching.

How to store?

- Store at room temperature and avoid sunlight
- After opening, it can be used until the expired date.
- After first opening the pouch, the medicinal product should be used within 6 months.